

WEST Search History

[Hide Items](#)[Restore](#)[Clear](#)[Cancel](#)

DATE: Wednesday, March 07, 2007

Hide?	<u>Set</u> <u>Name</u>	<u>Query</u>	<u>Hit</u> <u>Count</u>
		<i>DB=EPAB,JPAB,DWPI; PLUR=YES; OP=OR</i>	
<input type="checkbox"/>	L2	(luteinizing adj hormone adj receptor or lhr) and (hcg or human adj chorionic adj gonadotropin) and exoloop	0
		<i>DB=PGPB,USPT; PLUR=YES; OP=OR</i>	
<input type="checkbox"/>	L1	(luteinizing adj hormone adj receptor or lhr) and (hcg or human adj chorionic adj gonadotropin) and exoloop	2

END OF SEARCH HISTORY

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:ssptacmb1647

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

* * * * * Welcome to STN International * * * * *

NEWS	1		Web Page URLs for STN Seminar Schedule - N. America
NEWS	2		"Ask CAS" for self-help around the clock
NEWS	3	OCT 23	The Derwent World Patents Index suite of databases on STN has been enhanced and reloaded
NEWS	4	OCT 30	CHEMLIST enhanced with new search and display field
NEWS	5	NOV 03	JAPIO enhanced with IPC 8 features and functionality
NEWS	6	NOV 10	CA/CAPLUS F-Term thesaurus enhanced
NEWS	7	NOV 10	STN Express with Discover! free maintenance release Version 8.01c now available
NEWS	8	NOV 20	CA/CAPLUS to MARPAT accession number crossover limit increased to 50,000
NEWS	9	DEC 01	CAS REGISTRY updated with new ambiguity codes
NEWS	10	DEC 11	CAS REGISTRY chemical nomenclature enhanced
NEWS	11	DEC 14	WPIDS/WPINDEX/WPIX manual codes updated
NEWS	12	DEC 14	GBFULL and FRFULL enhanced with IPC 8 features and functionality
NEWS	13	DEC 18	CA/CAPLUS pre-1967 chemical substance index entries enhanced with preparation role
NEWS	14	DEC 18	CA/CAPLUS patent kind codes updated
NEWS	15	DEC 18	MARPAT to CA/CAPLUS accession number crossover limit increased to 50,000
NEWS	16	DEC 18	MEDLINE updated in preparation for 2007 reload
NEWS	17	DEC 27	CA/CAPLUS enhanced with more pre-1907 records
NEWS	18	JAN 08	CHEMLIST enhanced with New Zealand Inventory of Chemicals
NEWS	19	JAN 16	CA/CAPLUS Company Name Thesaurus enhanced and reloaded
NEWS	20	JAN 16	IPC version 2007.01 thesaurus available on STN
NEWS	21	JAN 16	WPIDS/WPINDEX/WPIX enhanced with IPC 8 reclassification data
NEWS	22	JAN 22	CA/CAPLUS updated with revised CAS roles
NEWS	23	JAN 22	CA/CAPLUS enhanced with patent applications from India
NEWS	24	JAN 29	PHAR reloaded with new search and display fields
NEWS	25	JAN 29	CAS Registry Number crossover limit increased to 300,000 in multiple databases
NEWS	26	FEB 13	CASREACT coverage to be extended
NEWS	27	Feb 15	PATDPASPC enhanced with Drug Approval numbers
NEWS	28	Feb 15	RUSSIAPAT enhanced with pre-1994 records
NEWS	29	Feb 23	KOREAPAT enhanced with IPC 8 features and functionality
NEWS	30	Feb 26	MEDLINE reloaded with enhancements
NEWS	31	Feb 26	EMBASE enhanced with Clinical Trial Number field
NEWS	32	Feb 26	TOXCENTER enhanced with reloaded MEDLINE
NEWS	33	Feb 26	IFICDB/IFIPAT/IFIUDB reloaded with enhancements
NEWS	34	Feb 26	CAS Registry Number crossover limit increased from 10,000 to 300,000 in multiple databases
NEWS EXPRESS			NOVEMBER 10 CURRENT WINDOWS VERSION IS V8.01c, CURRENT MACINTOSH VERSION IS V6.0c(ENG) AND V6.0Jc(JP), AND CURRENT DISCOVER FILE IS DATED 25 SEPTEMBER 2006.
NEWS HOURS			STN Operating Hours Plus Help Desk Availability

NEWS LOGIN Welcome Banner and News Items
NEWS IPC8 For general information regarding STN implementation of IPC 8
NEWS X25 X.25 communication option no longer available

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation of commercial gateways or other similar uses is prohibited and may result in loss of user privileges and other penalties.

* * * * * STN Columbus * * * * *

FILE 'HOME' ENTERED AT 14:16:52 ON 07 MAR 2007

=> file medline embase biosis caplus

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	0.21	0.21

FILE 'MEDLINE' ENTERED AT 14:17:01 ON 07 MAR 2007

FILE 'EMBASE' ENTERED AT 14:17:01 ON 07 MAR 2007
Copyright (c) 2007 Elsevier B.V. All rights reserved.

FILE 'BIOSIS' ENTERED AT 14:17:01 ON 07 MAR 2007
Copyright (c) 2007 The Thomson Corporation

FILE 'CAPLUS' ENTERED AT 14:17:01 ON 07 MAR 2007
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2007 AMERICAN CHEMICAL SOCIETY (ACS)

=> s (luteinizing hormone receptor or lhr) and (hcg or human chorionic gonadotropin) and exoloop

L1 11 (LUTEINIZING HORMONE RECEPTOR OR LHR) AND (HCG OR HUMAN CHORIONIC GONADOTROPIN) AND EXOLOOP

=> dup rem l1

PROCESSING COMPLETED FOR L1

L2 5 DUP REM L1 (6 DUPLICATES REMOVED)

=> dis his

(FILE 'HOME' ENTERED AT 14:16:52 ON 07 MAR 2007)

FILE 'MEDLINE, EMBASE, BIOSIS, CAPLUS' ENTERED AT 14:17:01 ON 07 MAR 2007

L1 11 S (LUTEINIZING HORMONE RECEPTOR OR LHR) AND (HCG OR HUMAN CHORIONIC GONADOTROPIN) AND EXOLOOP
L2 5 DUP REM L1 (6 DUPLICATES REMOVED)

=> dis ibib abs l2 1-5

L2	ANSWER 1 OF 5	MEDLINE on STN	DUPLICATE 1
ACCESSION NUMBER:	2001269988	MEDLINE	
DOCUMENT NUMBER:	PubMed ID: 10964934		
TITLE:	The role of the hinge region of the luteinizing hormone receptor in hormone interaction and signal generation.		
AUTHOR:	Zeng H; Phang T; Song Y S; Ji I; Ji T H		
CORPORATE SOURCE:	Department of Chemistry, University of Kentucky, Lexington, Kentucky 40506-0055, USA.		
CONTRACT NUMBER:	DK-51469 (NIDDK)		

HD-18702 (NICHD)
SOURCE: The Journal of biological chemistry, (2001 Feb 2) Vol. 276,
No. 5, pp. 3451-8. Electronic Publication: 2000-08-29.
Journal code: 2985121R. ISSN: 0021-9258.
PUB. COUNTRY: United States
DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)
(RESEARCH SUPPORT, U.S. GOV'T, P.H.S.)
LANGUAGE: English
FILE SEGMENT: Priority Journals
ENTRY MONTH: 200106
ENTRY DATE: Entered STN: 25 Jun 2001
Last Updated on STN: 5 Jan 2003
Entered Medline: 21 Jun 2001

AB Luteinizing hormone receptor, a G
protein-coupled receptor, consists of two halves, the N-terminal
extracellular hormone binding domain (exodomain) and the C-terminal
membrane-associated, signal-generating domain (endodomain). The exodomain
has seven to nine Leu-rich repeats, which are generally thought to form a
1/3 donut-like structure and interact with human choriogonadotropin (hCG).
The resulting hCG-exodomain complex adjusts the structure and its association
with the endodomain, which results in signal generation in the endodomain.
It is unclear whether the rigid 1/3 donut structure could provide the agility
and versatility of this dynamic action. In addition, there is no clue as to
where the endodomain contact point (the signal modulator) in the exodomain is.
To address these issues, the exodomain was examined by Ala scan and multiple
substitutions, while receptor peptides were used for photoaffinity labeling
and affinity cross-linking. Our results show that the C-flanking sequence
(hinge region), Thr(250)-Gln(268), of the Leu-rich repeats (LRRs) specifically
interacts with hCG, preferentially hCG α . This interaction is inhibited by
exoloop 2 of the endodomain but not by exoloops 1 and 3, suggesting an intimate
relationship between Thr(250)-Gln(268), exoloop 2, and hCG. Taken together,
our observations in this article suggest a new paradigm that the LRRs contact
the front of hCG, while both flanking regions of the LRRs interact with the
sides of hCG. This would trap hCG in the 1/3 donut structure of the LRRs
and enhance the binding affinity. In addition, mutations of conserved
Ser(255) in the sequence can constitutively activate the receptor. This
provides a clue for the signal modulator in the exodomain. In contrast,
a phenyl or phenolic group is necessary at conserved Tyr(253) for targeting
the receptor to the surface.

L2 ANSWER 2 OF 5 EMBASE COPYRIGHT (c) 2007 Elsevier B.V. All rights reserved on STN

ACCESSION NUMBER: 2003466657 EMBASE
TITLE: Hormone interactions to Leu-rich repeats in the gonadotropin receptors. III. Photoaffinity labeling of human chorionic gonadotropin with receptor Leu-rich repeat 4 peptide.
AUTHOR: Jeoung M.; Phang T.; Song Y.S.; Ji I.; Ji T.H.
CORPORATE SOURCE: T.H. Ji, Dept. of Chemistry, University of Kentucky, Lexington, KY 40506-0055, United States. tji@pop.uky.edu
SOURCE: Journal of Biological Chemistry, (2 Feb 2001) Vol. 276, No. 5, pp. 3443-3450. .
Refs: 34
ISSN: 0021-9258 CODEN: JBCHA3
COUNTRY: United States
DOCUMENT TYPE: Journal; Article
FILE SEGMENT: 003 Endocrinology
029 Clinical Biochemistry
LANGUAGE: English
SUMMARY LANGUAGE: English
ENTRY DATE: Entered STN: 30 Dec 2003

Last Updated on STN: 30 Dec 2003

AB Human chorionic gonadotropin (hCG) binds to the extracellular N-terminal domain, exodomain, of its receptor, and the resulting hCG-exodomain complex is thought to modulate the membrane associated domain, endodomain, of the receptor to generate hormone signal. The bulk of the exodomain is speculated to assume a crescent structure consisting of eight to nine Leu-rich repeats (LRRs), which may provide the hormone contact sites. Unfortunately, little experimental evidence is available for the precise hormone contact points in the exodomain and the endodomain. The two preceding articles (Song, Y., Ji, I., Beauchamp, J., Isaacs, N., and Ji, T. (2001) J. Biol. Chemical 276, 3426-3435; Song, Y., Ji, I., Beauchamp, J., Isaacs, N., and Ji, T. (2001) J. Biol. Chemical 276, 3436-3442) show that putative LRR2 and LRR4 are crucial for hormone binding. In particular, the N-terminal region of LRR4 assumes the hydrophobic core of the LRR4 loop, whereas the C-terminal region is crucial for signal generation. However, it is unclear whether LRR4 interacts hCG and the endodomain and how it might be involved in signal generation. In this article, our affinity labeling results present the first evidence that the N-terminal region of LRR4 interacts with hCG, preferentially the hCG α subunit and that the hCG/LRR4 complex interacts with exoloop 2 of the endodomain. This interaction offers a mechanism to generate hormone signal.

L2 ANSWER 3 OF 5 MEDLINE on STN DUPLICATE 2
ACCESSION NUMBER: 1999002809 MEDLINE
DOCUMENT NUMBER: PubMed ID: 9788760
TITLE: Surface retention of an inactivating lutropin receptor mutant in exoloop 3.
AUTHOR: Bhowmick N; Narayan P; Puett D
CORPORATE SOURCE: Department of Biochemistry and Molecular Biology, University of Georgia, Athens 30602, USA.
CONTRACT NUMBER: DK33973 (NIDDK)
SOURCE: Molecular and cellular biochemistry, (1998 Oct) Vol. 187, No. 1-2, pp. 221-7.
Journal code: 0364456. ISSN: 0300-8177.
PUB. COUNTRY: Netherlands
DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)
(RESEARCH SUPPORT, U.S. GOV'T, P.H.S.)
LANGUAGE: English
FILE SEGMENT: Priority Journals
ENTRY MONTH: 199812
ENTRY DATE: Entered STN: 15 Jan 1999
Last Updated on STN: 15 Jan 1999
Entered Medline: 18 Dec 1998

AB The heptahelical lutropin receptor (LHR) signals primarily via the Gs-adenylyl cyclase pathway and undergoes ligand-mediated receptor desensitization and internalization. A loss-of-function rat LHR mutant was recently described in which a single amino acid residue replacement in exoloop 3, K583E, had no effect on human choriongonadotropin (hCG) binding but essentially abolished signaling. This LHR mutant is a prime candidate for which to study hCG-mediated receptor internalization since it is highly unlikely that an amino acid residue in exoloop 3, i.e. an extracellular portion of LHR connecting transmembrane helices 6 and 7, could have any direct interaction with Galpha(s), which is located on the cytoplasmic face of the plasma membrane. A method to study endocytosis was adapted that involves concanavalin A binding to the glycoproteins on the cell surface, thus facilitating separation of the plasma membrane fraction from other cellular membrane fractions by sucrose gradient centrifugation. Conditions were used such that a single round of endocytosis could be determined with [125I]hCG. Endocytic rate constants of 0.03 and 0 min⁻¹ were obtained for LHR and the mutant, respectively, in transfected human embryonic kidney 293 cells;

moreover, internalization of the mutant could not be restored by the addition of 8-Br-cAMP. Thus, the presence of the second messenger cAMP is not sufficient for internalization of ligand-occupied LHR. Rather, it appears that ligand-mediated activation and subsequent internalization of LHR results from an altered conformational state or a conformation-dependent post-ligand binding modification such as phosphorylation.

L2 ANSWER 4 OF 5 EMBASE COPYRIGHT (c) 2007 Elsevier B.V. All rights reserved on STN

ACCESSION NUMBER: 96104114 EMBASE
DOCUMENT NUMBER: 1996104114
TITLE: Exoloop 3 of the luteinizing hormone/choriogonadotropin receptor. Lys583 is essential and irreplaceable for human choriogonadotropin (hCG)-dependent receptor activation but not for high affinity hCG binding.
AUTHOR: Ryu K.-S.; Gilchrist R.L.; Ji I.; Kim S.-J.; Ji T.H.
CORPORATE SOURCE: Dept. of Molecular Biology, University of Wyoming, Laramie, WY 82071-3944, United States
SOURCE: Journal of Biological Chemistry, (1996) Vol. 271, No. 13, pp. 7301-7304. .
ISSN: 0021-9258 CODEN: JBCHA3
COUNTRY: United States
DOCUMENT TYPE: Journal; Article
FILE SEGMENT: 029 Clinical Biochemistry
LANGUAGE: English
SUMMARY LANGUAGE: English
ENTRY DATE: Entered STN: 30 Apr 1996
Last Updated on STN: 30 Apr 1996

AB The luteinizing hormone/choriogonadotropin (CG) receptor belongs to a subfamily of glycoprotein hormone receptors within the seven-transmembrane receptor family. It is comprised of an extracellular N-terminal half of 341 amino acids and a membrane-associated C-terminal half of 303 amino acids. The N-terminal half is capable of high affinity hormone binding whereas the C-terminal half is capable of low affinity hormone binding and receptor activation. However, the precise location of the receptor activation site is currently unknown. We present evidence for the first time that Lys583 of exoloop 3 is crucial and irreplaceable for receptor activation to induce cAMP synthesis. Exoloop 3 is comprised of 11 amino acids and flanked by two Lys residues, Lys573 and Lys583, that are located at the boundaries with the transmembrane columns 6 and 7, respectively. All substitutions including Arg for Lys583 did not affect the high affinity human CG binding, but they resulted in the complete loss of cAMP synthesis induced by human CG. Ala substitutions of the other amino acids in exoloop 3 did not make such a dramatic impact on cAMP induction. The Ala scan revealed two distinct groups of amino acids in terms of their importance in cAMP induction, one group being more important than the other. Interestingly, these two groups of amino acids are arranged in an alternate sequence. This result suggests a specific structure similar to a β -like structure for exoloop 3.

L2 ANSWER 5 OF 5 EMBASE COPYRIGHT (c) 2007 Elsevier B.V. All rights reserved on STN

ACCESSION NUMBER: 97026945 EMBASE
DOCUMENT NUMBER: 1997026945
TITLE: Molecular mechanism of LH/CG receptor activation.
AUTHOR: Ryu K.-S.; Ji I.; Chang L.; Ji T.H.
CORPORATE SOURCE: T.H. Ji, Department of Molecular Biology, University of Wyoming, Laramie, WY 82071-3944, United States
SOURCE: Molecular and Cellular Endocrinology, (1996) Vol. 125, No. 1-2, pp. 93-100. .
Refs: 34

ISSN: 0303-7207 CODEN: MCEND6
 PUBLISHER IDENT.: S 0303-7207(96)03951-2
 COUNTRY: Ireland
 DOCUMENT TYPE: Journal; Conference Article
 FILE SEGMENT: 003 Endocrinology
 029 Clinical Biochemistry
 LANGUAGE: English
 SUMMARY LANGUAGE: English
 ENTRY DATE: Entered STN: 15 Feb 1997
 Last Updated on STN: 15 Feb 1997

AB It is known that the N-terminal half of the LH/CG receptor is responsible for high hCG binding whereas the C-terminal half is capable of receptor activation. Our results suggest that initial hCG binding at the high affinity site in the N-half receptor induces conformational adjustments. This leads to low affinity secondary contacts of the complex of hCG/the N-half receptor with the C-half receptor. This low affinity secondary contact is responsible for activating the receptor. This is based on the following observations. The C-terminal tail of hCG α is known to be involved in activation of the LH/CG receptor. In addition to hCG, we examined the C-terminal three residues (His90-Lys91-Ser92) of the common α subunit of FSH and TSH. The results show their differential roles in the three hormones. Ser92 is important for binding and cAMP induction of TSH but not for hCG and FSH. Lys91 is important for binding and cAMP induction of hCG, and cAMP induction but not binding of FSH. It is not important for binding or cAMP induction of TSH. His90 is important for all three hormones. When all three residues were truncated, FSH and TSH lose their affinity for binding and cAMP induction, whereas hCG is still capable of binding but not cAMP induction. Therefore, the three amino acids contribute differently in receptor binding and cAMP induction of hCG, FSH and TSH. Our data also indicate that the evolution of the α subunit has been constrained in order not to impair any one of the hormones. This suggests that each hormone can be independently engineered to improve the potency. To chemically identify the contact site of the α C-tail of hCG in the LH/CG receptor, a decamer peptide corresponding to the α subunit sequence from His83 to Ser92 (peptide α 83-92) was derivatized with UV sensitive reagent, ABG and radio-iodinated. The resulting ABG-125I-peptide α 83-92 was capable of binding and activating the LH/CG receptor. Furthermore, it specifically photoaffinity-labeled the LH/CG receptor. In addition, the amino group of α Lys91 of peptide α 83-92 is crosslinked to a carboxyl group of the receptor, an indication of close association. Reciprocal mutagenesis of α Lys91 and Asp397 in exoloop 1 of the LH/CG receptor suggests the complementary of this pair in receptor activation but not the high affinity interaction of hCG and the receptor. In addition, Lys583 of exoloop 3 is also crucial for receptor activation. To test the conformational adjustment, ABG was attached to hCG α and reassociated with untreated β to produce ABG-125I- α/β . The extent of inter-subunit crosslinking of ABG-125I- α/β bound to the receptor was two to three fold less than unbound ABG-125I- α/β . This result indicates structural change at the subunit interface in response to hCG binding to the receptor.

=> FIL STNGUIDE
 COST IN U.S. DOLLARS
 FULL ESTIMATED COST

SINCE FILE	TOTAL
ENTRY	SESSION
31.69	31.90

FILE 'STNGUIDE' ENTERED AT 14:19:14 ON 07 MAR 2007
 USE IS SUBJECT TO THE TERMS OF YOUR CUSTOMER AGREEMENT
 COPYRIGHT (C) 2007 AMERICAN CHEMICAL SOCIETY, JAPAN SCIENCE

AND TECHNOLOGY CORPORATION, AND FACHINFORMATIONSZENTRUM KARLSRUHE

FILE CONTAINS CURRENT INFORMATION.

LAST RELOADED: Mar 2, 2007 (20070302/UP).

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:ssptacmb1647

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

* * * * * Welcome to STN International * * * * *

NEWS	1		Web Page URLs for STN Seminar Schedule - N. America
NEWS	2		"Ask CAS" for self-help around the clock
NEWS	3	OCT 23	The Derwent World Patents Index suite of databases on STN has been enhanced and reloaded
NEWS	4	OCT 30	CHEMLIST enhanced with new search and display field
NEWS	5	NOV 03	JAPIO enhanced with IPC 8 features and functionality
NEWS	6	NOV 10	CA/CAPLUS F-Term thesaurus enhanced
NEWS	7	NOV 10	STN Express with Discover! free maintenance release Version 8.01c now available
NEWS	8	NOV 20	CA/CAPLUS to MARPAT accession number crossover limit increased to 50,000
NEWS	9	DEC 01	CAS REGISTRY updated with new ambiguity codes
NEWS	10	DEC 11	CAS REGISTRY chemical nomenclature enhanced
NEWS	11	DEC 14	WPIDS/WPINDEX/WPIX manual codes updated
NEWS	12	DEC 14	GBFULL and FRFULL enhanced with IPC 8 features and functionality
NEWS	13	DEC 18	CA/CAPLUS pre-1967 chemical substance index entries enhanced with preparation role
NEWS	14	DEC 18	CA/CAPLUS patent kind codes updated
NEWS	15	DEC 18	MARPAT to CA/CAPLUS accession number crossover limit increased to 50,000
NEWS	16	DEC 18	MEDLINE updated in preparation for 2007 reload
NEWS	17	DEC 27	CA/CAPLUS enhanced with more pre-1907 records
NEWS	18	JAN 08	CHEMLIST enhanced with New Zealand Inventory of Chemicals
NEWS	19	JAN 16	CA/CAPLUS Company Name Thesaurus enhanced and reloaded
NEWS	20	JAN 16	IPC version 2007.01 thesaurus available on STN
NEWS	21	JAN 16	WPIDS/WPINDEX/WPIX enhanced with IPC 8 reclassification data
NEWS	22	JAN 22	CA/CAPLUS updated with revised CAS roles
NEWS	23	JAN 22	CA/CAPLUS enhanced with patent applications from India
NEWS	24	JAN 29	PHAR reloaded with new search and display fields
NEWS	25	JAN 29	CAS Registry Number crossover limit increased to 300,000 in multiple databases
NEWS	26	FEB 13	CASREACT coverage to be extended
NEWS	27	Feb 15	PATDPASPC enhanced with Drug Approval numbers
NEWS	28	Feb 15	RUSSIAPAT enhanced with pre-1994 records
NEWS	29	Feb 23	KOREAPAT enhanced with IPC 8 features and functionality
NEWS	30	Feb 26	MEDLINE reloaded with enhancements
NEWS	31	Feb 26	EMBASE enhanced with Clinical Trial Number field
NEWS	32	Feb 26	TOXCENTER enhanced with reloaded MEDLINE
NEWS	33	Feb 26	IFICDB/IFIPAT/IFIUDB reloaded with enhancements
NEWS	34	Feb 26	CAS Registry Number crossover limit increased from 10,000 to 300,000 in multiple databases

NEWS EXPRESS NOVEMBER 10 CURRENT WINDOWS VERSION IS V8.01c, CURRENT MACINTOSH VERSION IS V6.0c(ENG) AND V6.0Jc(JP), AND CURRENT DISCOVER FILE IS DATED 25 SEPTEMBER 2006.

NEWS HOURS STN Operating Hours Plus Help Desk Availability

NEWS LOGIN Welcome Banner and News Items
NEWS IPC8 For general information regarding STN implementation of IPC 8
NEWS X25 X.25 communication option no longer available

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation of commercial gateways or other similar uses is prohibited and may result in loss of user privileges and other penalties.

* * * * * STN Columbus * * * * *

FILE 'HOME' ENTERED AT 12:44:08 ON 07 MAR 2007

=> file medline embase biosis caplus
COST IN U.S. DOLLARS

SINCE FILE	TOTAL
ENTRY	SESSION
0.21	0.21

FULL ESTIMATED COST

FILE 'MEDLINE' ENTERED AT 12:44:52 ON 07 MAR 2007

FILE 'EMBASE' ENTERED AT 12:44:52 ON 07 MAR 2007
Copyright (c) 2007 Elsevier B.V. All rights reserved.

FILE 'BIOSIS' ENTERED AT 12:44:52 ON 07 MAR 2007
Copyright (c) 2007 The Thomson Corporation

FILE 'CAPLUS' ENTERED AT 12:44:52 ON 07 MAR 2007
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2007 AMERICAN CHEMICAL SOCIETY (ACS)

=> s contraception and (LHR or luteinizing(w)hormone(w)receptor) and
(leu(w)rich(w)repeats or lrr or exodomain(w)1 or exodomain(w)2 or exodomain(w)3)
L1 0 CONTRACEPTION AND (LHR OR LUTEINIZING(W) HORMONE(W) RECEPTOR)
AND (LEU(W) RICH(W) REPEATS OR LRR OR EXODOMAIN(W) 1 OR EXODOMAI
N(W) 2 OR EXODOMAIN(W) 3)

=> s contraception and (leu(w)rich(w)repeats or lrr or exodomain(w)1 or
exodomain(w)2 or exodomain(w)3)
L2 0 CONTRACEPTION AND (LEU(W) RICH(W) REPEATS OR LRR OR EXODOMAIN(W)
1 OR EXODOMAIN(W) 2 OR EXODOMAIN(W) 3)

=> dis his

(FILE 'HOME' ENTERED AT 12:44:08 ON 07 MAR 2007)

FILE 'MEDLINE, EMBASE, BIOSIS, CAPLUS' ENTERED AT 12:44:52 ON 07 MAR 2007
L1 0 S CONTRACEPTION AND (LHR OR LUTEINIZING(W)HORMONE(W)RECEPTOR) A
L2 0 S CONTRACEPTION AND (LEU(W)RICH(W)REPEATS OR LRR OR EXODOMAIN(W)